



Office de la propriété
intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Commission

An Agency of
Industry Canada

PCT / CA 00/00770

0/019890
20 JUL 2000 (20.07.00)

4

*Bureau canadien
des brevets
Certification*

*Canadian Patent
Office
Certification CA00/770*

La présente atteste que les documents
ci-joints, dont la liste figure ci-dessous,
sont des copies authentiques des docu-
ments déposés au Bureau des brevets.

This is to certify that the documents
attached hereto and identified below are
true copies of the documents on file in
the Patent Office.

Specification as originally filed with Application for Patent Serial No: 2,276,798, on June
30, 1999, by KEN CURRY, XINYAO DU AND HASAN PAJOUHESH, for
"2-Aminoindane Analogs"

REC'D 01 AUG 2000

AIPO

PCT

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

S. Lachance
Agent certificateur/Certifying Officer

July 20, 2000

Date

Canada

(CIPO 68)

OPIC



CIPO

2-AMINOINDANE ANALOGS

ABSTRACT

The present invention relates to therapeutically active (4,9-Benzyl)-aminoindane compounds, a method of preparing the same, and to pharmaceutical compositions comprising the compounds. The novel compounds are useful in treating diseases of the central nervous system related to the metabotropic glutamate receptor system.

FIELD OF THE INVENTION

This invention pertains to therapeutically active (4,9-Benzyl)-aminoindane derivatives, a method for preparing the same, pharmaceutical compositions comprising the compounds and a method of treating diseases of the Central Nervous System (CNS) therewith.

BACKGROUND OF THE INVENTION

The acidic amino acid L-Glutamate is recognized as the major excitatory neurotransmitter in the CNS. The receptors that respond to L-Glutamate are called excitatory amino acid receptors. The excitatory amino acid receptors are thus of great physiological importance, playing a role in a variety of physiological processes, such as long-term potentiation (learning and memory), the development of synaptic plasticity, motor control, respiratory and cardiovascular regulation, and sensory perception.

Excitatory amino acid receptors are classified into two general types and both are activated by L-Glutamic acid and its analogs. Receptors activated by L-Glutamic acid that are directly coupled to the opening of cation channels in the cell membrane of the neurons are termed "ionotropic." This type of receptor has been subdivided into at least three subtypes, which are defined by the depolarizing actions of the selective agonists N-Methyl-D-aspartate (NMDA), α -Amino-3-hydroxy-5-methylisoxazole-4-propionic acid (AMPA), and Kainic acid (KA).

The second general type of receptor is the G-protein or second messenger-linked "metabotropic" excitatory amino acid receptor. This second type is coupled to multiple second messenger systems that lead to enhanced phosphoinositide hydrolysis, activation of phospholipase D, increases or decreases in cAMP formation, and changes in ion channel function (Schoepp and Conn, *Trends in Pharmacological Science*, 14:13, 1993). Both types of receptors appear not only to mediate normal synaptic transmission along excitatory pathways but also to participate in the modification of synaptic connections during development and throughout life.

So far eight different clones of the G-protein-coupled mGluRs have been identified (Knopfel et al., 1995, *J. Med. Chem.*, 38, 1417-1426). These receptors function to modulate the presynaptic release of L-Glutamate, and the postsynaptic sensitivity of the neuronal cell to L-Glutamate excitation. Based on pharmacology, sequence homology and the signal transduction pathway that they activate, the mGluRs have been subclassified into three groups. The mGluR₁ and mGluR₅ receptors form group I. They are coupled to hydrolysis of phosphatidylinositol (PI) and are selectively activated by (R,S)-3,5-dihydroxyphenylglycine (Brabet et al., *Neuropharmacology*, 34, 895-903, 1995). Group II comprises mGluR₂ and mGluR₃ receptors. They are negatively coupled to adenylate cyclase and are selectively activated by (2S,1'R,2'R,3'R)-2-(2,3-dicarboxycyclopropyl)glycine (DCG-IV; Hayashi et al., *Nature*, 366, 687-690, 1993). Finally, the mGluR₄, mGluR₆, mGluR₇ and mGluR₈ receptors belong to group III. They are also negatively coupled to adenylate cyclase and are selectively activated by (L)-2-amino-4-phosphonobutyric acid (L-AP4; Knopfel et al., 1995, *J. Med. Chem.*, 38, 1417-1426).

Agonists and antagonists of these receptors are believed useful for the treatment of acute and chronic neurodegenerative conditions, and as antipsychotic, anticonvulsant, analgesic, anxiolytic, antidepressant, and anti-emetic agents. Antagonists and agonists of neural receptors are classified as selective for a particular receptor or receptor subtype, or as non-selective. Antagonists may also be classified as competitive or non-competitive. While competitive and non-competitive antagonists act on the receptors in a different manner to produce similar results, selectivity is based upon the observations that some antagonists exhibit high levels of activity at a single receptor type, and little or no activity at other receptors. In the case of receptor-specific diseases and conditions, the selective agonists and antagonists are of the most value.

Compounds such as L-Glutamic acid, Quisqualic acid and Ibotenic acid are known to act as non-selective agonists on the mGluRs, while selective ionotropic glutamate receptor agonists such

as NMDA, AMPA and kainate have little effect on these receptors. Recently a few compounds without activity at the ionotropic glutamate receptors but with activity at the metabotropic receptors have been identified. These include *trans*-ACPD (*trans*-(1*S*,3*R*)-1-aminocyclopentane-1,3-dicarboxylic acid), the partial agonist *L*-AP3 (*L*-2-amino-3-phosphonopropionic acid; Palmer, E., Monaghan, D. T. and Cotman, C. W. *Eur. J. Pharmacol.* 166, 585-587, 1989; Desai, M. A. and Conn, P. J. *Neuroscience Lett.* 109, 157-162, 1990; Schoepp, D. D. et al., *J. Neurochemistry* 56, 1789-1796, 1991; Schoepp D. D. and Johnson B. G. *J. Neurochemistry* 53, 1865-1913, 1989), *L*-AP4 (*L*-2-amino-4-phosphonobutyric acid) which is an agonist at the mGluR₄ receptor (Thomsen C. et al., *Eur. J. Pharmacol.* 227, 361-362, 1992) and some of the isomers of CCG (2-(carboxycyclopropyl)glycines) especially *L*-CCG-I and *L*-CCG-II (Hayashi, Y. et al., *Br. J. Pharmacol.* 107, 539-543, 1992).

Very few selective antagonists at the mGluRs have been reported. However some phenylglycine derivatives, *S*-4CPG (*S*-4-carboxyphenylglycine), *S*-4C3HPG (*S*-4-carboxy-3-hydroxyphenylglycine) and *S*-MCPG (*S*- α -methyl-4-carboxyphenylglycine) have been reported to antagonize *trans*-ACPD-stimulated phosphoinositide hydrolysis and thus possibly act as antagonists at mGluR₁ and mGluR₅ subtypes (Thomsen, C. and Suzdak, P., *Eur. J. Pharmacol.* 245, 299, 1993).

Research directed towards mGluRs is beginning to show that mGluRs may be implicated in a number of normal as well as pathological mechanisms in the brain and spinal cord. For example, activation of these receptors on neurons can: influence levels of alertness, attention and cognition; protect nerve cells from excitotoxic damage resulting from ischemia, hypoglycemia and anoxia; modulate the level of neuronal excitation; influence central mechanisms involved in controlling movement; reduce sensitivity to pain; reduce levels of anxiety.

The use of compounds active at the mGluRs for the treatment of epilepsy is corroborated by investigations of the influence of *trans*-ACPD on the formation of convulsions (Sacaan and Schoepp, *Neuroscience Lett.* 139, 77, 1992) and that phosphoinositide hydrolysis mediated via mGluR is increased after kindling experiments in rats (Akiyama et al. *Brain Res.* 569, 71, 1992). *Trans*-ACPD has been shown to increase release of dopamine in the rat brain, which indicates that compounds acting on the mGluRs might be usable for the treatment of Parkinson's disease and Huntington's Chorea (Sacaan et al., *J. Neurochemistry* 59, 245, 1992).

Trans-ACPD has also been shown to be a neuroprotective agent in a medial cerebral artery occlusion (MCAO) model in mice (Chiamulera et al. *Eur. J. Pharmacol.* 215, 353, 1992), and it has been shown to inhibit NMDA-induced neurotoxicity in nerve cell cultures (Koh et al., *Proc. Natl. Acad. Sci. USA* 88, 9431, 1991). The mGluR-active compounds are also implicated in the treatment of pain. This is proved by the fact that antagonists at the mGluRs antagonize sensory synaptic response to noxious stimuli of thalamic neurons (Eaton, S. A. et al., *Eur. J. Neuroscience*, 5, 186, 1993).

The use of compounds active at the mGluRs for treatment of neurological diseases such as senile dementia have also been indicated by the findings of Zheng and Gallagher, *Neuron* 9, 163, 1992 and Bashir et al. (*Nature* 363, 347, 1993) who demonstrated that activation of mGluRs is necessary for the induction of long-term potentiation (LTP) in nerve cells (septal nucleus, hippocampus) and the finding that long-term depression is induced after activation of mGluRs in cerebellar granule cells (Linden et al. *Neuron* 7, 81, 1991).


Thus compounds that demonstrate either activating or inhibiting activity at mGluRs have therapeutic potential for the treatment of neurological disorders. These compounds have application as new drugs to treat both acute and chronic neurological disorders, such as stroke and head injuries; epilepsy; movement disorders associated with Parkinson's disease and Huntington's chorea; pain; anxiety; AIDS dementia; and Alzheimer's disease. Since the mGluRs can influence levels of alertness, attention and cognition; protect nerve cells from excitotoxic damage resulting from ischemia, hypoglycemia and anoxia; modulate the level of neuronal excitation; influence central mechanisms involved in controlling movement; reduce sensitivity to pain; and reduce levels of anxiety, these compounds can also be used to influence these situations and also find use in learning and memory deficiencies such as senile dementia. mGluRs may also be involved in addictive behavior, alcoholism, drug addiction, sensitization and drug withdrawal (*Science*, 280:2045, 1998), so compounds acting at mGluRs might also be used to treat these disorders.

The current pharmaceutical options for treating neurological disorders tend to be very general and non-specific in their actions in that, although they may reduce the clinical symptoms associated with a specific neurological disorder, they may also negatively impact normal function of the central nervous system of patients. Thus new cellular targets and drugs that are more specific in

their actions require to be identified and developed and thus a need remains for chemical compounds that demonstrate specific binding characteristics towards mGluRs.

SUMMARY OF THE INVENTION

It is an object of this invention to provide novel compounds that demonstrate activity at the various metabotropic glutamate receptors. In particular, a compound of Formula I and stereoisomers thereof:



wherein:

R1, and **R2** must independently be either:

- 1) H; or:
- 2) an acidic group selected from the group consisting of carboxy, phosphono, phosphino, sulfono, sulfino, borono, tetrazol, isoxazol, $-(CH_2)_n$ -carboxy, $-(CH_2)_n$ -phosphono, $-(CH_2)_n$ -phosphino, $-(CH_2)_n$ -sulfono, $-(CH_2)_n$ -sulfino, $-(CH_2)_n$ -borono, $-(CH_2)_n$ -tetrazol, and $-(CH_2)_n$ -isoxazol, where $n = 1, 2, 3, 4, 5$, or 6 ; or:

X can be an acidic group selected from the group consisting of carboxy, phosphono, phosphino, sulfono, sulfino, borono, tetrazol, isoxazol.

Y can be a basic group selected from the group consisting of 1° amino, 2° amino, 3° amino, quaternary ammonium salts, aliphatic 1° amino, aliphatic 2° amino, aliphatic 3° amino, aliphatic quaternary ammonium salts, aromatic 1° amino, aromatic 2° amino, aromatic 3° amino, aromatic quaternary ammonium salts, imidazol, guanidino, boronoamino, allyl, urea, thiourea ;

m can be 0, 1.

R3, R4, R5, R6 are independently H, nitro, amino, halogen, tritium, trifluoromethyl, trifluoroacetyl, sulfo, carboxy, carbamoyl, sulfamoyl or acceptable esters thereof;

or a salt thereof with a pharmaceutically acceptable acid or base.

DETAILED DESCRIPTION OF THE INVENTION

The terms and abbreviations used in the instant examples have their normal meanings unless otherwise designated. For example "°C" refers to degrees Celsius; "N" refers to normal or normality; "mmol" refers to millimole or millimoles; "g" refers to gram or grams; "ml" means milliliter or milliliters; "M" refers to molar or molarity; "p-" refers to para, "MS" refers to mass spectrometry; "IR" refers to infrared spectroscopy; and "NMR" refers to nuclear magnetic resonance spectroscopy.


As would be understood by the skilled artisan, throughout the synthesis of the compounds of Formula I it may be necessary to employ an amino-protecting group or a carboxy-protecting group in order to reversibly preserve a reactively susceptible amino or carboxy functionality while reacting other functional groups on the compound.

Examples of such amino-protecting groups include formyl, trityl, phthalimido, trichloroacetyl, chloroacetyl, bromoacetyl, iodoacetyl, and urethane-type blocking groups such as benzyloxycarbonyl, 4-phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl, 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxycarbonyl, *t*-butoxycarbonyl, 2-(4-xenyl)-isopropoxycarbonyl, 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(*p*-toluyl)-prop-2-yloxycarbonyl, cyclopentanyloxy-carbonyl, 1-methylcyclopentanyloxy-carbonyl, cyclohexanyloxy-carbonyl, 1-methylcyclohexanyloxy-carbonyl, 2-methylcyclohexanyloxy-carbonyl, 2-(4-toluylsulfonyl)-ethoxycarbonyl, 2-(methylsulfonyl)ethoxycarbonyl, 2-(triphenylphosphino)-ethoxycarbonyl, fluorenylmethoxycarbonyl ("FMOC"),

2-(trimethylsilyl)ethoxycarbonyl, allyloxycarbonyl,
 1-(trimethylsilylmethyl)prop-1-enyloxycarbonyl, 5-benzisoxalylmethoxycarbonyl,
 4-acetoxymethyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-propoxycarbonyl,
 cyclopropylmethoxycarbonyl, 4-(decyloxy)benzyloxycarbonyl, isobornyloxycarbonyl,
 1-piperidyloxycarbonyl and the like; benzoylmethylsulfonyl group, 2-nitrophenylsulfonyl,
 diphenylphosphine oxide and like amino-protecting groups. The species of amino-protecting
 group employed is not critical so long as the derivatized amino group is stable to the condition of
 subsequent reaction(s) on other positions of the intermediate molecule and can be selectively
 removed at the appropriate point without disrupting the remainder of the molecule including any
 other amino-protecting group(s). Preferred amino-protecting groups are *t*-butoxycarbonyl
 (*t*-Boc), allyloxycarbonyl and benzyloxycarbonyl (CbZ). Further examples of these groups are
 found in E. Haslam, *Protecting Groups in Organic Chemistry*, (J. G. W. McOmie, ed., 1973), at
 Chapter 2; and T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, (1991),
 at Chapter 7.

Examples of such carboxy-protecting groups include methyl, *p*-nitrobenzyl, *p*-methylbenzyl,
p-methoxybenzyl, 3,4-dimethoxybenzyl, 2,4-dimethoxybenzyl, 2,4,6-trimethoxybenzyl,
 2,4,6-trimethylbenzyl, pentamethylbenzyl, 3,4-methylenedioxybenzyl, benzhydryl,
 4,4'-dimethoxybenzhydryl, 2,2',4,4'-tetramethoxybenzhydryl, *t*-butyl, *t*-amyl, trityl,
 4-methoxytrityl, 4,4'-dimethoxytrityl, 4,4',4''-trimethoxytrityl, 2-phenylprop-2-yl, trimethylsilyl,
t-butyldimethylsilyl, phenacyl, 2,2,2-trichloroethyl, β -(di(*n*-butyl)methylsilyl)ethyl,
p-toluenesulfonylethyl, 4-nitrobenzylsulfonylethyl, allyl, cinnamyl,
 1-(trimethylsilylmethyl)prop-1-en-3-yl and like moieties. Preferred carboxy-protecting groups are
 allyl, benzyl and *t*-butyl. Further examples of these groups are found in E. Haslam, *supra*, at
 Chapter 5; and T. W. Greene and P. G. M. Wuts, *supra*, at Chapter 5.

The present invention provides a compound of the formula I and stereoisomers thereof:



R1, and **R2** must independently be either :

1) H; or:

2) an acidic group selected from the group consisting of carboxy, phosphono, phosphino, sulfono, sulfino, borono, tetrazol, isoxazol, $-(CH_2)_n$ -carboxy, $-(CH_2)_n$ -phosphono, $-(CH_2)_n$ -phosphino, $-(CH_2)_n$ -sulfono, $-(CH_2)_n$ -sulfino, $-(CH_2)_n$ -borono, $-(CH_2)_n$ -tetrazol, and $-(CH_2)_n$ -isoxazol, where $n = 1, 2, 3, 4, 5$, or 6 ; or

X can be an acidic group selected from the group consisting of carboxy, phosphono, phosphino, sulfono, sulfino, borono, tetrazol, isoxazol.

Y can be a basic group selected from the group consisting of 1° amino, 2° amino, 3° amino, quaternary ammonium salts, aliphatic 1° amino, aliphatic 2° amino, aliphatic 3° amino, aliphatic quaternary ammonium salts, aromatic 1° amino, aromatic 2° amino, aromatic 3° amino, aromatic quaternary ammonium salts, imidazol, guanidino, boronoamino, allyl, urea, thiourea ;

m can be 0, 1.

R3, **R4**, **R5**, **R6** are independently H, nitro, amino, halogen, tritium, trifluoromethyl, trifluoroacetyl, sulfo, carboxy, carbamoyl, sulfamoyl or acceptable esters thereof;

or a salt thereof with a pharmaceutically acceptable acid or base.

In particular compounds wherein the compound of Formula I is selected from the group consisting of:

R1 = CO_2H , CH_2CO_2H

R2 = H

X = CO_2H and **Y** = NH_2

R1 = H

R2 = CO_2H , CH_2CO_2H

X = CO_2H and **Y** = NH_2

While all of the compounds of Formula I are believed to demonstrate activity at the metabotropic glutamate receptors (mGluRs), certain groups of Formula I compounds are more preferred for such use.

As noted supra, this invention includes the pharmaceutically acceptable salts of the compounds defined by Formula I. A compound of this invention can possess a sufficiently acidic, a sufficiently basic, or both functional groups, and accordingly react with any of a number of organic and inorganic bases, and inorganic and organic acids, to form a pharmaceutically acceptable salt.

The term "pharmaceutically acceptable salt" as used herein, refers to salts of the compounds of the above formula which are substantially non-toxic to living organisms. Typical pharmaceutically acceptable salts include those salts prepared by reaction of the compounds of the present invention with a pharmaceutically acceptable mineral or organic acid or an organic or inorganic base. Such salts are known as acid addition and base addition salts.

Acids commonly employed to form acid addition salts are inorganic acids such as hydrochloric acid, hydrobromic acid, hydriodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids such as *p*-toluenesulfonic acid, methanesulfonic acid, oxalic acid, *p*-bromophenylsulfonic acid, carbonic acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like. Examples of such pharmaceutically acceptable salts are the sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, hydrochloride, dihydrochloride, isobutyrate, caproate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, hydroxybenzoate, methoxybenzoate, phthalate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, γ -hydroxybutyrate, glycolate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate and the like.

Preferred pharmaceutically acceptable acid addition salts are those formed with mineral acids such as hydrochloric acid and hydrobromic acid, and those formed with organic acids such as maleic acid and methanesulfonic acid.

Salts of amine groups may also comprise quarternary ammonium salts in which the amino nitrogen carries a suitable organic group such as an alkyl, alkenyl, alkynyl, or aralkyl moiety.

Base addition salts include those derived from inorganic bases, such as ammonium or alkali or alkaline earth metal hydroxides, carbonates, bicarbonates, and the like. Such bases useful in preparing the salts of this invention thus include sodium hydroxide, potassium hydroxide, ammonium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, potassium bicarbonate, calcium hydroxide, calcium carbonate, and the like. The potassium and sodium salt forms are particularly preferred.

It should be recognized that the particular counterion forming a part of any salt of this invention is usually not of a critical nature, as long as the salt as a whole is pharmacologically acceptable and as long as the counterion does not contribute undesired qualities to the salt as a whole.

This invention further encompasses the pharmaceutically acceptable solvates of the compounds of Formula I. Many of the Formula I compounds can combine with solvents such as water, methanol, ethanol and acetonitrile to form pharmaceutically acceptable solvates such as the corresponding hydrate, methanolate, ethanolate and acetonitrilate.

The compounds of the present invention have multiple asymmetric (chiral) centers. As a consequence of these chiral centers, the compounds of the present invention occur as racemates, mixtures of enantiomers and as individual enantiomers, as well as diastereomers and mixtures of diastereomers. All asymmetric forms, individual isomers and combinations thereof, are within the scope of the present invention.

The prefixes "*R*" and "*S*" are used herein as commonly used in organic chemistry to denote the absolute configuration of a chiral center, according to the Cahn-Ingold-Prelog system. The stereochemical descriptor *R* (*rectus*) refers to that configuration of a chiral center with a clockwise relationship of groups tracing the path from highest to second-lowest priorities when viewed from the side opposite to that of the lowest priority group. The stereochemical descriptor *S* (*sinister*) refers to that configuration of a chiral center with a counterclockwise relationship of groups tracing the path from highest to second-lowest priority when viewed from the side opposite to the lowest priority group. The priority of groups is decided using sequence rules as described by Cahn et al., *Angew. Chem.*, 78, 413-447, 1966 and Prelog, V. and Helmchen, G.; *Angew. Chem. Int. Ed. Eng.*, 21, 567-583, 1982).


In addition to the *R,S* system used to designate the absolute configuration of a chiral center, the older D-L system is also used in this document to denote relative configuration, especially with reference to amino acids and amino acid derivatives. In this system a Fischer projection of the compound is oriented so that carbon-1 of the parent chain is at the top. The prefix "D" is used to represent the relative configuration of the isomer in which the functional (determining) group is on the right side of the carbon atom at the chiral center and "L", that of the isomer in which it is on the left.

As would be expected, the stereochemistry of the Formula I compounds is critical to their potency as agonists. The relative stereochemistry is preferably established early during synthesis, which avoids stereoisomer separation problems later in the process. Subsequent synthetic steps then employ stereospecific procedures so as to maintain the preferred configuration. The preferred methods of this invention are the methods employing those preferred compounds.

Non-toxic metabolically labile esters and amides of compounds of Formula I are ester or amide derivatives of compounds of Formula I that are hydrolyzed in vivo to afford said compounds of Formula I and a pharmaceutically acceptable alcohol or amine. Examples of metabolically labile esters include esters formed with (1-6C) alkanols in which the alkanol moiety may be optionally substituted by a (1-8C) alkoxy group, for example methanol, ethanol, propanol and methoxyethanol. Examples of metabolically labile amides include amides formed with amines such as methylamine.


According to another aspect, the present invention provides a process for the preparation of a compound of Formula I, or a pharmaceutically acceptable metabolically labile ester or amide thereof, or a pharmaceutically acceptable salt thereof, which comprises:

(a) hydrolyzing a compound of formula II :




in which **R1**, **R2**, **R3**, **R4**, **R5** and **R6** have the meaning defined above. **R7** represents a hydrogen atom or an acyl group. Preferred values for **R7** are hydrogen and (2-6C) alkanoyl groups, such as acetyl.

(b) hydrolyzing a compound of formula III:



in which **R1**, **R2**, **R3**, **R4**, **R5** and **R6** have the meaning defined above, **R8** and **R9** each independently represent a hydrogen atom, a (2-6C) alkanoyl group, a (1-4C) alkyl group, a (3-4C) alkenyl group or a phenyl (1-4C) alkyl group in which the phenyl is unsubstituted or substituted by halogen, (1-4C) alkyl or (1-4C) alkoxy, or a salt thereof; or:

(c) deprotecting a compound of formula IV:



in which **R1**, **R2**, **R3**, **R4**, **R5** and **R6** have the meaning defined above and **R10** represents a

hydrogen atom or a carboxyl protecting group, or a salt thereof, and R11 represents a hydrogen atom or a nitrogen protecting group;

whereafter, if necessary and/or desired:

- (i) resolving the compound of Formula I;
- (ii) converting the compound of Formula I into a non-toxic metabolically labile ester or amide thereof; and/or;
- (iii) converting the compound of Formula I or a non-toxic metabolically labile ester or amide thereof into a pharmaceutically acceptable salt thereof.

The protection of carboxylic acid and amine groups is generally described in McOmie, *Protecting Groups in Organic Chemistry*, Plenum Press, NY, 1973, and Greene and Wuts, *Protective Groups in Organic Synthesis*, 2nd. Ed., John Wiley & Sons, NY, 1991. Examples of carboxy protecting groups include alkyl groups such as methyl, ethyl, *t*-butyl and *t*-amyl; aralkyl groups such as benzyl, 4-nitrobenzyl, 4-methoxybenzyl, 3,4-dimethoxybenzyl, 2,4-dimethoxybenzyl, 2,4,6-trimethoxybenzyl, 2,4,6-trimethylbenzyl, benzhydryl and trityl; silyl groups such as trimethylsilyl and *t*-butyldimethylsilyl; and allyl groups such as allyl and 1-(trimethylsilylmethyl)prop-1-en-3-yl. Examples of amine protecting groups include acyl groups, such as groups of formula R11 CO in which R11 represents (1-6C) alkyl, (3-10C) cycloalkyl, phenyl(1-6C) alkyl, phenyl(1-6C) alkoxy, or a (3-10C) cycloalkoxy, wherein a phenyl group may optionally be substituted by one or two substituents independently selected from amino, hydroxy, nitro, halogeno, (1-6C) alkyl, (1-6C) alkoxy, carboxy, (1-6C) alkoxycarbonyl, carbamoyl, (1-6C) alkanoylamino, (1-6C) alkylsulphonylamino, phenylsulphonylamino, toluenesulphonylamino, and (1-6C) fluoroalkyl.


The compounds of Formula II are conveniently hydrolyzed in the presence of an acid, such as hydrochloric acid or sulfuric acid, or a base, such as an alkali metal hydroxide, for example sodium hydroxide. The hydrolysis is conveniently performed in an aqueous solvent such as water and at a temperature in the range from 50° to 200° C.

The compounds of Formula III are conveniently hydrolyzed in the presence of a base, for example an alkali metal hydroxide such as lithium, sodium or potassium hydroxide, or an alkaline earth

metal hydroxide such as barium hydroxide. Suitable reaction media include water. The temperature is conveniently in the range from 50° to 150° C.

The compounds of Formula IV may be deprotected by conventional methods. Thus, an alkyl carboxyl protecting group may be removed by hydrolysis. The hydrolysis may conveniently be performed by heating the compound of Formula IV in the presence of either a base, for example an alkali metal hydroxide such as lithium, sodium or potassium hydroxide, or an alkaline metal hydroxide, such as barium hydroxide, or an acid such as hydrochloric acid. The hydrolysis is conveniently performed at a temperature in the range from 10° to 300° C. An aralkyl carboxyl-protecting group may conveniently be removed by hydrogenolysis. The hydrogenolysis may conveniently be effected by reacting the compound of Formula IV with hydrogen in the presence of a Group VIII metal catalyst, for example a palladium catalyst such as palladium on charcoal. Suitable solvents for the reaction include alcohols such as ethanol. The reaction is conveniently performed at a temperature in the range from 0° to 100°C. An acyl amine protecting group is also conveniently removed by hydrolysis, for example as described for the removal of an alkyl carboxyl protecting group.


The compounds of Formula II may be prepared by reacting a compound of formula V:



in which R1, R2, R3, R4, R5 and R6 have the meaning defined above, with an alkali metal cyanide, such as lithium, sodium or potassium cyanide, and either ammonium carbonate in an aqueous alcohol, such as aqueous ethanol, or with an ammonium halide, such as ammonium chloride, conveniently in the presence of ultrasound. If the reaction is conducted with ammonium carbonate, the reaction is conveniently performed at a temperature in the range from 35° to 150°C. If desired, the compounds of Formula II may then be alkylated, for example using a

compound of formula RCl. As described in more detail hereinafter, the alkylated compounds may be readily separated into their diastereomers. If the reaction is conducted with an ammonium halide in the presence of ultrasound, the ammonium halide is mixed with chromatography grade alumina in the presence of a suitable diluent such as acetonitrile. The mixture is then irradiated with ultrasound, whereafter the compound of Formula V is added, and the mixture is again irradiated. The alkali metal cyanide is then added, followed by further irradiation with ultrasound.


Individual isomers of compounds of Formula II may be made by reacting a compound of the Formula V with the stereoisomers of the chiral agent (S)- and (R)-phenylglycinol and a reactive cyanide such as trimethylsilyl cyanide to form the intermediate compounds of Formula VI.



The compounds of Formula III may be prepared by reacting a compound of Formula V with an alkali metal cyanide, such as lithium, sodium or potassium cyanide, and ammonium carbonate or ammonium carbamate. Common solvents include alcohols, such as methanol, aqueous methanol and aqueous ethanol. Conveniently the reaction is performed at a temperature in the range of from 10° to 150 °C. If desired, the compounds of Formula III may then be alkylated, for example using an appropriate alkyl, aryl or acyl chloride.


The compounds of Formula (V) may be prepared from compounds of Formula VII, either by hydrolysis or by alkylation followed by hydrolysis of the resulting compounds.

Where R can be any carboxyl-protecting group as previously described on page 7.




VII

Compounds of Formula VII are prepared from compounds of Formula VIII by Dieckmann cyclization:




VIII

Compounds of Formula VIII may be prepared by esterification of compounds of Formula IX:



IX

Compounds of Formula IX may be formed from commercially available 1,2-phenylenediacetonitrile (X) by hydrolysis in aqueous HCl.



X

Many of the intermediates described herein, for example the compounds of Formula II, III and IV are believed to be novel, and are provided as further aspects of the invention.

The Formula I compounds of the present invention are agonists or antagonists at certain metabotropic glutamate receptors (mGluRs). Therefore, another aspect of the present invention is a method of affecting mGluRs in mammals, which comprises administering to a mammal requiring modulated excitatory amino acid neurotransmission a pharmacologically-effective amount of a compound of Formula I. The term "pharmacologically-effective amount" is used to represent an amount of the compound of the invention that is capable of affecting the mGluRs. By affecting, a compound of the invention is acting as an agonist or antagonist. When a compound of the invention acts as an agonist, the interaction of the compound with the excitatory amino acid receptor mimics the response of the interaction of this receptor with its natural ligand, L-Glutamic acid. When a compound of the invention acts as an antagonist, the interaction of the compound with the excitatory amino acid receptor blocks the response of the interaction of this receptor with its natural ligand, L-Glutamic acid.

The particular dose of compound administered according to this invention will, of course, be determined by the particular circumstances surrounding the case, including the compound administered, the route of administration, the particular condition being treated, and similar considerations. The compounds can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, or intranasal routes. Alternatively, the compound may be administered by continuous infusion. A typical daily dose will contain from about 0.001 mg/kg to about 100 mg/kg of the active compound of this invention. Preferably, daily doses will be about 0.05 mg/kg to about 50 mg/kg, more preferably from about 0.1 mg/kg to about 20 mg/kg.

A variety of physiological functions have been shown to be subject to influence by excessive or inappropriate stimulation of excitatory amino acid transmission. The Formula I compounds of the present invention are believed (through their interactions at the mGluRs) to have the ability to treat a variety of neurological disorders in a warm-blooded mammal associated with this condition, including acute neurological disorders such as cerebral deficits subsequent to cardiac bypass surgery and grafting, cerebral ischemia (e.g. stroke and cardiac arrest), spinal cord trauma, head trauma, perinatal hypoxia, and hypoglycemic neuronal damage. The Formula I compounds are believed to have the ability to treat a variety of chronic neurological disorders, such as Alzheimer's disease, Huntington's Chorea, amyotrophic lateral sclerosis, AIDS-induced dementia, ocular damage and retinopathy, cognitive disorders, and idiopathic and drug-induced Parkinson's disease. The present invention also provides methods for treating these disorders which comprises administering to a patient in need thereof an effective amount of a compound of Formula I.

The Formula I compounds of the present invention (through their interactions at the mGluRs) are also believed to have the ability to treat a variety of other neurological disorders in mammals that are associated with glutamate dysfunction, including muscular spasms, convulsions, migraine headaches, urinary incontinence, psychosis, drug tolerance, withdrawal, and cessation (i.e. opiates, benzodiazepines, nicotine, cocaine, or ethanol), smoking cessation, anxiety and related disorders (e.g. panic attack), emesis, brain edema, chronic pain, sleep disorders, Tourette's syndrome, attention deficit disorder, and tardive dyskinesia. Therefore, the present invention also provides methods for treating these disorders which comprise administering to a patient in need thereof an effective amount of the compound of Formula I.

The Formula I compounds of the present invention (through their interactions at the mGluRs) are also believed to have the ability to treat a variety of psychiatric disorders, such as schizophrenia, anxiety and related disorders (e.g. panic attack), depression, bipolar disorders, psychosis, and obsessive compulsive disorders. The present invention also provides methods for treating these disorders which comprises administering to a patient in need thereof an effective amount of a compound of Formula I.

The pharmacological properties of the compounds of the invention can be illustrated by determining their effects in various functional in vitro assays. The compounds of the invention

were studied in an in vitro assay that measured the effect on PI hydrolysis or the formation of cyclic AMP in Chinese hamster ovary cell lines expressing mGluR_{1α}, mGluR₂ and mGluR_{4a} cloned metabotropic glutamate receptors.

Principle

So far eight different clones of the G-protein-coupled mGluRs have been identified (Knopfel et al., 1995, *J. Med. Chem.*, 38, 1417-1426). These receptors function to modulate the presynaptic release of L-Glutamic acid, and the postsynaptic sensitivity of the neuronal cell to L-Glutamic acid excitation. Based on pharmacology, sequence homology and the signal transduction pathway that they activate, the mGluRs have been subclassified into three groups. The mGluR₁ and mGluR₅ receptors form group I. They are coupled to hydrolysis of phosphatidylinositol (PI) and are selectively activated by (RS)-3,5-dihydroxyphenylglycine (Brabet et al., *Neuropharmacology*, 34, 895-903, 1995). Group II comprises mGluR₂ and mGluR₃ receptors. They are negatively coupled to adenylate cyclase and are selectively activated by (2S,1'R,2'R,3'R)-2-(2,3-dicarboxycyclopropyl)glycine (DCG-IV; Hayashi et al., *Nature*, 366, 687-690, 1993). Finally, the mGluR₄, mGluR₆, mGluR₇ and mGluR₈ receptors belong to group III. They are also negatively coupled to adenylate cyclase and are selectively activated by (L)-2-amino-4-phosphonylbutyric acid (L-AP4; Knopfel et al., 1995, *J. Med. Chem.*, 38, 1417-1426).

Cell Culture


The Chinese hamster ovary cell lines expressing mGluR_{1α}, mGluR₂ and mGluR_{4a} receptors have been described previously (Aramori and Nakanishi, *Neuron* 8, 757-765; 1992; Tanabe et al., *Neuron* 8, 169-179, 1992; Tanabe et al., *J. Neurosci.* 13, 1372-1378). They were maintained at 37°C in a humidified 5% CO₂ incubator in Dulbecco's Modified Eagle Medium (DMEM) containing a reduced concentration of (S)-glutamine (2mM) and were supplemented with 1% proline, penicillin (100 U/ml), streptomycin (100 mg/ml) and 10% dialyzed fetal calf serum (all GIBCO, Paisley). Two days before assay 1.8×10^6 cells were divided into the wells of 24 well plates.

Second Messenger Assays

PI hydrolysis was measured as described previously (Hayashi et al., *Br. J. Pharmacol.* 107, 539-543, 1992; Hayashi et al., *J. Neurosci.* 14, 3370-3377, 1994). Briefly, the cells were labeled with [^3H]inositol (2 $\mu\text{Ci/ml}$) 24 h prior to the assay. For agonist assays, the cells were incubated with ligand dissolved in phosphate-buffered saline (PBS)-LiCl for 20 min, and agonist activity was determined by measurement of the level of ^3H -labeled mono-, bis- and tris-inositol phosphates by ion-exchange chromatography. For antagonist assays, the cells were preincubated with the ligand dissolved in PBS-LiCl for 20 min prior to incubation with ligand and 10 μM (L)-Glutamic acid for 20 min. The antagonist activity was then determined as the inhibitory effect of the (L)-Glutamic acid mediated response. The assay of cyclic AMP formation was performed as described previously (Hayashi et al., *Br. J. Pharmacol.* 107, 539-543, 1992; Hayashi et al., *J. Neurosci.* 14, 3370-3377, 1994). Briefly, the cells were incubated for 10 min in PBS containing the ligand and 10 μM forskolin and 1mM 3-Isobutyl-1-methylxanthine (IBMX; both Sigma, St. Louis, MO, USA). The agonist activity was then determined as the inhibitory effect of the Forskolin-induced cyclic AMP formation. For antagonist assay, the cells were preincubated with ligand dissolved in PBS containing 1 mM IBMX for 20 min prior to a 10 min incubation in PBS containing the ligand, 20 μM (mGluR₂) or 50 μM (mgluR_{4a}), (L)-Glutamic acid, 10 μM Forskolin and 1 mM IBMX.

Results

Some of the compounds of the invention were tested for antagonist activity against Chinese hamster ovary cell lines expressing mGluR_{1a}, mGluR₂ and mGluR_{4a} cloned mGluRs at a concentration of 5 mM. When tested as antagonists of the increase in PI hydrolysis evoked by 10 μM (L)-Glutamic acid, some compounds of the invention effectively blocked this increase in PI hydrolysis by an action at the mGluR_{1a} receptor. The data for two of the compounds of the invention are shown in Figure 1 below.



According to another aspect, the present invention provides a method of modulating one or more mGluR functions in a warm-blooded mammal which comprises administering an effective amount of a compound of Formula I, or a non-toxic metabolically-labile ester or amide thereof, or a pharmaceutically acceptable salt thereof.

The compounds of the present invention are preferably formulated prior to administration. Therefore, another aspect of the present invention is a pharmaceutical formulation comprising a compound of Formula I and a pharmaceutically-acceptable carrier, diluent, or excipient. The present pharmaceutical formulations are prepared by known procedures using well-known and readily available ingredients. In making the compositions of the present invention, the active ingredient will usually be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier, and may be in the form of a capsule, sachet, paper, or other container. When the carrier serves as a diluent, it may be a solid, semi-solid, or liquid material that acts as a vehicle, excipient, or medium for the active ingredient.

The compounds of Formula I are usually administered in the form of pharmaceutical compositions. These compounds can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, and intranasal. These compounds are effective as both injectable and oral compositions. Such compositions are prepared in a manner well known in the pharmaceutical art and comprise at least one active compound.

The present invention also provides pharmaceutical compositions containing compounds as disclosed in the claims in combination with one or more pharmaceutically acceptable, inert or physiologically active, diluents or adjuvants. The compounds of the invention can be freeze-dried and, if desired, combined with other pharmaceutically acceptable excipients to prepare formulations for administration. These compositions may be presented in any form appropriate for the administration route envisaged. The parenteral and the intravenous route are the preferential routes for administration.

Compounds of the general Formula I may be administered orally, topically, parenterally, by inhalation or spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein

includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition, there is provided a pharmaceutical formulation comprising a compound of general Formula I and a pharmaceutically acceptable carrier. One or more compounds of general Formula I may be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or adjuvants and, if desired, other active ingredients. The pharmaceutical compositions containing compounds of general Formula I may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs.

Compositions intended for oral use may be prepared according to any known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavouring agents, colouring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients that are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methyl cellulose, hydropropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example hepta-decaethyleneoxycetanol, or

condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or *n*-propyl- *p*-hydroxy benzoate, one or more colouring agents, one or more flavouring agents or one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredients in a vegetable oil, for example peanut oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavouring agents may be added to provide palatable oral preparations. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavouring and colouring agents, may also be present.

Pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oil phase may be a vegetable oil, for example olive oil or peanut oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavouring and colouring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to

known art using those suitable dispersing or wetting agents and suspending agents that have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or a suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compound(s) of the general Formula I may be administered, together or separately, in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

Compound(s) of general Formula I may be administered, together or separately, parenterally in sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anaesthetics, preservatives and buffering agents can be dissolved in the vehicle.

The dosage to be administered is not subject to defined limits, but it will usually be an effective amount. It will usually be the equivalent, on a molar basis of the pharmacologically active free form produced from a dosage formulation upon the metabolic release of the active free drug to achieve its desired pharmacological and physiological effects. The compositions are preferably formulated in a unit dosage form, each dosage containing from about 0.05 to about 100 mg, more usually about 1.0 to about 30 mg, of the active ingredient. The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

The active compound is effective over a wide dosage range. For examples, dosages per day normally fall within the range of about 0.01 to about 30 mg/kg of body weight. A typical daily dose will contain from about 0.01 mg/kg to about 100 mg/kg of the active compound of this invention. Preferably, daily doses will be about 0.05 mg/kg to about 50 mg/kg, more preferably from about 0.1 mg/kg to about 25 mg/kg. In the treatment of adult humans, the range of about 0.1

to about 15 mg/kg/day, in single or divided dose, is especially preferred. However, it will be understood that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, and the severity of the patient's symptoms, and therefore the above dosage ranges are not intended to limit the scope of the invention in any way. In some instances dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect, provided that such larger doses are first divided into several smaller doses for administration throughout the day.

The compositions are preferably formulated in a unit dosage form, each dosage containing from about 5 mg to about 500 mg, more preferably about 25 mg to about 300 mg of the active ingredient. The term "unit dosage form" refers to a physically discrete unit suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical carrier, diluent, or excipient. The following formulation examples are illustrative only and are not intended to limit the scope of the invention in any way.

Formulation 1

Hard gelatin capsules are prepared using the following ingredients:

	Quantity (mg/capsule)
Active Ingredient	250
Starch, dried	200
Magnesium stearate	10
Total	460

The above ingredients are mixed and filled into hard gelatin capsules in 460 mg quantities.

Formulation 2

A tablet is prepared using the ingredients below:

Quantity (mg/tablet)

Active Ingredient	250
Cellulose, microcrystalline	400
Silicon dioxide, fumed	10
Stearic acid	5
Total	665

The components are blended and compressed to form tablets each weighing 665 mg.

Formulation 3

An aerosol solution is prepared containing the following components:

	Weight %
Active Ingredient	0.25
Ethanol	29.75
Propellant 22 (Chlorodifluoromethane)	70.00
Total	100

The active compound is mixed with ethanol and the mixture added to a portion of the Propellant 22, cooled to -30 °C and transferred to a filling device. The required amount is then fed to a stainless steel container and diluted with the remainder of the propellant. The valve units are then fitted to the container.

Formulation 4

Tablets each containing 60 mg of active ingredient are made as follows:

Quantity (mg/tablet)

Active Ingredient	60
Starch	45
Microcrystalline cellulose	35
Polyvinylpyrrolidone	4
Sodium carboxymethyl starch	4.5
Magnesium stearate	0.5
Talc	1.0
Total	150

The active ingredient, starch, and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders that are then passed through a No. 14 mesh U.S. sieve. The granules so produced are dried at 50°C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate, and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

Formulation 5

Capsules each containing 80 mg medicament are made as follows:

	Quantity (mg/capsule)
Active Ingredient	80
Starch	59
Microcrystalline cellulose	59
Magnesium stearate	2
Total	200

The active ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 45 sieve, and filled into hard gelatin capsules in 200 mg quantities.

Formulation 6

Suppositories each containing 225 mg of active ingredient may be made as follows:

	Quantity (mg/suppository)
Active Ingredient	225
Saturated fatty acid glycerides	2000
Total	2225

The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2 g capacity and allowed to cool.

Formulation 7

Suspensions each containing 50 mg of medicament per 5 mL dose are made as follows:

Active Ingredient	50 mg
Sodium carboxymethyl cellulose	50 mg
Syrup	1.25 mL
Benzoic acid solution	0.10 mL
Flavour	q.v.
Color	q.v.
Purified water to total	5 mL

The medicament is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor and color are diluted with some of the water and added, with stirring. Sufficient water is then added to produce the required volume.

Formulation 8

An intravenous formulation may be prepared as follows:

	Quantity
Active Ingredient	100 mg
Mannitol	100 mg
5 N Sodium hydroxide	200 mL
Purified water to total	5 mL

Formulation 9

A topical formulation may be prepared as follows:

	Quantity
Active Ingredient	1-10 g
Emulsifying Wax	30 g
Liquid Paraffin	20 g
White soft paraffin to	100 g

The white soft paraffin is heated until molten. The liquid paraffin and emulsifying wax are incorporated and stirred until dissolved. The active ingredient is added and stirring is continued until dispersed. The mixture is then cooled until solid.

Formulation 10

Sublingual or buccal tablets, each containing 10 mg of active ingredient, may be prepared as follows:

	Quantity (mg/tablet)
Active Ingredient	10.0
Glycerol	210.5

Water	143.0
Sodium Citrate	4.5
Polyvinyl Alcohol	26.5
Polyvinylpyrrolidone	15.5
Total	410.0

The glycerol, water, sodium citrate, polyvinyl alcohol, and polyvinylpyrrolidone are admixed together by continuous stirring and maintaining the temperature at about 90 °C. When the polymers have gone into solution, the solution is cooled to about 50°-55 °C and the medicament is slowly admixed. The homogeneous mixture is poured into forms made of an inert material to produce a drug-containing diffusion matrix having a thickness of about 2-4 mm. This diffusion matrix is then cut to form individual tablets having the appropriate size.

Another preferred formulation employed in the methods of the present invention employs transdermal delivery devices ("patches"). Such transdermal patches may be used to provide continuous or discontinuous infusion of the compounds of the present invention in controlled amounts.


The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art (see, for example, U.S. Pat. No. 5,023,252, issued Jun. 11, 1991) herein incorporated by reference. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

Frequently, it will be desirable or necessary to introduce the pharmaceutical composition to the brain, either directly or indirectly. Direct techniques usually involve placement of a drug delivery catheter into the host's ventricular system to bypass the blood-brain barrier. One such implantable delivery system, used for the transport of biological factors to specific anatomical regions of the body, is described in U.S. Pat. No. 5,011,472, issued Apr. 30, 1991, which is herein incorporated by reference.

Indirect techniques, which are generally preferred, usually involve formulating the compositions to provide for drug latention by the conversion of hydrophilic drugs into lipid-soluble drugs or prodrugs. Latention is generally achieved through blocking of the hydroxy, carbonyl, sulfate, and primary amine groups present on the drug to render the drug more lipid soluble and amenable

to transportation across the blood-brain barrier. Alternatively, the delivery of hydrophilic drugs may be enhanced by intra-arterial infusion of hypertonic solutions that can transiently open the blood-brain barrier.

EXAMPLES

*Preparation 1: 1,2-phenylenediacetic acid*

In a round-bottomed flask, fitted with a reflux condenser, 10 g of phenylenediacetonitrile and 180 g of 50% sulphuric acid are refluxed for 2 h. The reaction mixture is cooled on ice to yield (2) as white crystals. Yield 11.4 g. This compound was used without further purification.

Preparation 2: Dimethyl-1,2-phenylenediacetate

The diacid (2) from above is dissolved in 150 mL of methanol, saturated with HCl gas and refluxed for 2 h. The resulting solution is cooled, evaporated to dryness and the residue taken up in 200 mL of diethyl ether. The ethereal extract is washed with saturated sodium bicarbonate solution, dried over magnesium sulphate and evaporated to give crude (3). The residue is purified by flash chromatography on silica to yield 11.1 g 78% from (1).

Preparation 3: Methyl-2-indanone-1-carboxylate

A solution of (3) (10 g) in benzene (50 mL) is placed in a round-bottomed flask together with molecular sodium (1.5 g) and stirred until no further hydrogen was given off. The resulting solution is washed with 1 N HCl (50 mL), water (50 mL) and brine (50 mL), dried over magnesium sulphate and evaporated to give crude (4) which was re-crystallized from water/methanol to give 8 g (92%) of (4) as a light yellow solid. NMR (CDCl₃), 3.18 (s, 2H, enolate), 3.59 (s, 2H), 3.79 (s, 3H, enolate), 3.97 (s, 3H), 7.05-7.62 (4H), 11.03 (s, 1H), 12.13 (s, 1H, enolate). The ratio of ketone:enolate 5.6:1.

Preparation 4: Ethyl-1-methoxycarbonyl-2-oxo-indaneacetate

A solution containing 6.0 g of (4) and 9.44 mL of DBU and 7.0 mL of ethylbromoacetate in 25 mL of DMF, is stirred overnight at r.t. The reaction mixture is diluted with ethyl acetate (50 mL) and washed with water, dried over magnesium sulphate and evaporated to give crude (5). The crude mixture is purified by flash chromatography (ethyl acetate:hexanes 10-30%:90-70%) to yield 7.1 g (oil) 82% of (5). NMR (CDCl₃), 3.20-3.42 (AB Pattern, 2H), 3.59 (s, 3H), 3.70-3.95 (4H), 7.15-7.35 (4H).

Preparation 5: 2-oxo-indaneacetic acid

A mixture of (5) and 50 mL of 1:1 6 N HCl:acetic acid is refluxed for 2 h. The resulting mixture gives (6) on cooling, which is re-crystallized from water/methanol to yield 3.5 g 71% of (6). NMR (CDCl₃) 2.90-3.10 (m, 2H), 3.58 (s, 2H), 3.70 (m, 1H), 7.1-7.4 (4H).

Preparation 6: 1-indaneacetic acid-2-(5,5'-hydantoin)


1.9 g of the keto acid (6) were dissolved in 20 mL of water in a sealed tube together with 1.3 g (2.0 mmol) of potassium cyanide and 3.8 g of ammonium carbonate (4.0 mmol). The solution is stirred and heated to 90-100°C for 16 h. The resulting dark solution is cooled and acidified with 6 N HCl and concentrated to a residue of (7) which is extracted into ethanol, filtered and used without further purification.

Preparation 7: 2-amino-2-carboxyindaneacetic acid

The hydantoin (7) was placed in a round-bottomed flask and refluxed together with 45 mL of 2 N NaOH for 16 h. The resulting solution is cooled, acidified with 6 N HCl, evaporated to dryness and extracted into ethanol. The ethanolic solution is treated with propylene oxide to precipitate a mixture of crude *cis* and *trans* (8). Chromatography on Spectrum 1X4 anion exchange resin, with dilute acetic acid gave *cis* and *trans* (8) as colorless crystals from methanol/water. *Cis* (8) ¹H NMR (D₂O) δ 2.5-2.7 (m, 2H), 3.15 (d, 1H), 3.58 (d, 1H), 4.05 (m, 1H), 7.1-7.35 (m, 4H). *Trans* (8) ¹H NMR (D₂O) δ 2.55 (dd, 1H), 2.8 (dd, 1H), 3.23 (d, 1H), 3.55 (d, 1H), 3.85 (dd, 1H), 7.15-7.3 (m, 4H).

We claim:

I. A compound of the formula:



Wherein **R1**, and **R2** can each separately be selected from the group consisting of:

- (i) H
- (ii) an acidic group selected from the group consisting of carboxy, phosphono, phosphino, sulfono, sulfinio, borono, tetrazol, isoxazol, $-(CH_2)_n$ -carboxy, $-(CH_2)_n$ -phosphono, $-(CH_2)_n$ -phosphino, $-(CH_2)_n$ -sulfono, $-(CH_2)_n$ -sulfinio, $-(CH_2)_n$ -borono, $-(CH_2)_n$ -tetrazol, and $-(CH_2)_n$ -isoxazol, where $n = 1, 2, 3, 4, 5$, or 6 ; or
X can be an acidic group selected from the group consisting of carboxy, phosphono, phosphino, sulfono, sulfinio, borono, tetrazol, isoxazol.
Y can be a basic group selected from the group consisting of 1° amino, 2° amino, 3° amino, quaternary ammonium salts, aliphatic 1° amino, aliphatic 2° amino, aliphatic 3° amino, aliphatic quaternary ammonium salts, aromatic 1° amino, aromatic 2° amino, aromatic 3° amino, aromatic quaternary ammonium salts, imidazol, guanidino, boronoamino, allyl, urea, thiourea ;
- (ii) a basic group selected from the group consisting of 1° amino, 2° amino, 3° amino, quaternary ammonium salts, aliphatic 1° amino, aliphatic 2° amino, aliphatic 3° amino, aliphatic quaternary ammonium salts, aromatic 1° amino, aromatic 2° amino, aromatic 3°


amino, aromatic quaternary ammonium salts, imidazol, guanidino, boronoamino, allyl, urea, thiourea

m can be 0, 1.

R4, R5, R6, R7 are independently H, nitro, amino, halogen, tritium, trifluoromethyl, trifluoroacetyl, sulfo, carboxy, carbamoyl, sulfamoyl, -or an acceptable ester thereof.


2. A compound as claimed in claim 1, wherein R1 can be H, CO₂H, CH₂CO₂H, XY, CH₂XY
3. A compound as claimed in claim 1, wherein R2 can be H, CO₂H, CH₂CO₂H, XY, CH₂XY
4. A process for the preparation of a compound of Formula I, or a pharmaceutically acceptable metabolically-labile ester or amide thereof, or a pharmaceutically acceptable salt thereof, which comprises:

(a) hydrolyzing a compound of formula:




In which R1, R2, R3, R4, R5 and R6 have the meaning defined above. R7 represents a hydrogen atom or an acyl group. Preferred values for R7 are hydrogen and (2-6C) alkanoyl groups, such as acetyl.

(b) hydrolyzing a compound of formula:



in which **R1**, **R2**, **R3**, **R4**, **R5** and **R6** have the meaning defined above, **R8** and **R9** each independently represent a hydrogen atom, a (2-6C) alkanoyl group, a (1-4C) alkyl group, a (3-4C) alkenyl group or a phenyl (1-4C) alkyl group in which the phenyl is unsubstituted or substituted by halogen, (1-4C) alkyl or (1-4C) alkoxy, or a salt thereof or:

(c) deprotecting a compound of formula:




in which **R1**, **R2**, **R3**, **R4**, **R5** and **R6** have the meaning defined above and **R10** represents a hydrogen atom or a carboxyl protecting group, or a salt thereof, and **R11** represents a hydrogen atom or a nitrogen protecting group;

whereafter, if necessary and/or desired:

(i) resolving the compound of Formula I;


- (ii) converting the compound of Formula I into a non-toxic metabolically labile ester or amide thereof and/or;
- (iii) converting the compound of Formula I or a non-toxic metabolically labile ester or amide thereof into a pharmaceutically acceptable salt thereof.

5. A pharmaceutical formulation, which comprises a compound as claimed in claim 1 and a pharmaceutically acceptable carrier, diluent or excipient.
6. A method of modulating one or more metabotropic glutamate receptor functions in a warm blooded mammal requiring such treatment, which comprises administering an effective amount of a compound as claimed in claim 1.
7. A compound of formula:




in which R1, R2, R3, R5, R5 and R6 have the meaning defined above. R7 represents a hydrogen atom or an acyl group. Preferred values for R7 are hydrogen and (2-6C) alkanoyl groups, such as acetyl.

8. A compound of formula:



in which **R1**, **R2**, **R3**, **R4**, **R5** and **R6** have the meaning defined above, **R8** and **R9** each independently represent a hydrogen atom, a (2-6C) alkanoyl group, a (1-4C) alkyl group, a (3-4C) alkenyl group or a phenyl (1-4C) alkyl group in which the phenyl is unsubstituted or substituted by halogen, (1-4C) alkyl or (1-4C) alkoxy, or a salt thereof or:

9. A compound of formula:



in which **R1**, **R2**, **R3**, **R4**, **R5** and **R6** have the meaning defined above and **R10** represents a hydrogen atom or a carboxyl protecting group, or a salt thereof, and **R11** represents a hydrogen atom or a nitrogen protecting group.

THIS PAGE BLANK (USPTO)

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)